

Station

Diagnosis and treatment of common diseases in children (pediatrics, neonatology, children's infectious diseases)

1. Professional standard

Integrated management of childhood illnesses

2. Competencies to be checked

Detection and assesment of general signs of danger.

Conclusion by the algorithm of the traffic light.

Determination of necessary actions.

Y. Diagonosis and treatment of local bacterial infection of the baby's umbilical wound

Check for signs of danger and other symptoms:

1. Redness of the umbilical wound.
2. Purulent discharge from umbilical wound.
3. Pustules on the skin
4. Make a conclusion according to the algorithm "traffic light".
5. Prescribe Amoxicillin Oral Suspension 125 mg / 5 ml 2 times per day for 5 days.
6. Teach mother how to treat a local infection at home: Wash your hands. Carefully wash the umbilical wound with boiled water. Dry the umbilical wound. Lay down Methylenum coeruleum solution in ethanol. Wash your hands.
7. Observe how mother treat the baby for the first time.
8. Inform the mother about conditions that require immediate re-treatment.
9. Re-examination In 2 days.
10. If there is any improvement, recommend the mother. Continue prescribed treatment for 5 days.
11. If the redness of the umbilical wound persists or purulent discharge appears, or the condition worsens: To direct the baby to the hospital.