

Algorithm of student actions

"Tooth extraction"

1. Say orally: "The patient is has an indication for tooth extraction 1.3"
2. Wear gloves
3. Treat hands with antiseptic
4. Say orally: "Please open your mouth"
5. Stand to the right of the phantom
6. Take a saliva ejector and secure it in the phantom's mouth
7. Clean the operating area with a dry napkin
8. Treat the operating area with a napkin basked in antiseptic
9. Say orally: "I will administer a right infraorbital anesthesia to the patient"
10. Say orally: "I will administer the patient incisional anesthesia"
11. Grasp the thumb and forefinger of the alveolar process of the tooth about to be extracted on the vestibular and palatal side
12. Take a trowel and perform a syndesmotomy (removal of the circular ligament of the tooth)
13. Choose straight forceps from the available tools
14. Apply forceps to the crown of the tooth so that the cheeks of the forceps are superimposed on the vestibular and palatal side of the tooth
15. Push the forceps to the CEJ of the tooth
16. Fix the tongs
17. Carry out luxation first in the vestibular side, then on the palatal side
18. Rotation
19. Carry out traction (removal of a tooth from a hole)
20. Place the tooth and forceps in the tray
21. Take a curettage spoon
22. Carry out an debridement of the socket (removal of pieces of alveoli and granulations)
23. Apply a gauze swab to the socket of the removed tooth
24. Remove the saliva ejector
25. Give recommendations to the patient on postoperative care of a wound in house conditions:

- do not rinse!
- do not place your tongue into the socket of the removed tooth
- Do not eat for 2 hours